

# Dermatology Order Form

Fax completed form to: \_\_\_\_\_



| PATIENT INFORMATION   |   |   |  |
|---|---|---|--|
| Patient Name:   | Date of Birth:  | Referral Date:  |  |
| Address:  |   | City/State/Zip:   |  |
| Home Phone:   | Cell Phone:   | Work Phone:   |  |
| Secondary Contact:  | Height:   | Weight:   | Male Female                              |
| Patient Diagnosis & ICD-10:   |   |   |  |
| Allergies:  |   |   |  |
| PROVIDER INFORMATION  |   |   |  |
| Physician Name:   | Lic.#:  | DEA #:  |  |
| Practice Name:  |   | NPI#:   |  |
| Address:  |   | City/State/Zip:   |  |
| Office Contact:   | Phone:  | Fax:  |  |
| Supervisory Physician (if applicable):  |   |   |  |
| PLEASE ATTACH   |   |   |  |
| Patient demographics & front/back copy of all insurance cards (prescription & medical)  |   | TB lab results within last 12 months ( <i>Stelara, Simponi Aria, Ilumya &amp; Infiximabs only</i> ) |  |
| Recent office visit notes, history & physical, lab & pertinent procedure results  |   | HBV lab results within last 12 months ( <i>Infiximabs &amp; Simponi Aria only</i> )                 |  |
| Current medication list & list of prior medications tried and failed (with dates)   |   | Letter of medical necessity if drug dosing or indication is outside of FDA guidelines               |  |
| NURSING & LAB ORDERS  |   |   |  |
| <b>Nurse Orders:</b> Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.                           |   |   |  |
| <b>Flush Orders:</b> NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line    |   |   |  |
| <b>Lab Orders:</b>  |   | <b>Lab Date &amp; Frequency:</b>  |  |
| PRESCRIPTION ORDERS   |   |   |  |
| <b>Anaphylaxis Kit:</b>   | Epinephrine 0.3mg IM as needed  | Solu-cortef 250mg-500mg IV as needed  | Solu-Medrol 60mg - 125mg IV as needed    |
| (Check all that apply)  | Diphenhydramine _____ mg IV as needed   | NS Hydration 500 ml IV over 30 minutes as needed  | Other _____                              |
| <b>Pre-Medications:</b>   | Acetaminophen _____ mg PO _____ minutes prior to infusion   | Solu-Medrol _____ mg IV _____ minutes prior to infusion   |  |
| (Check all that apply)  | Diphenhydramine _____ mg PO ---OR--- IV infusion _____ minutes prior to infusion  | Other _____   |  |
| <b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary   |   |   |  |
| PRODUCT   | PRESCRIPTION INFORMATION  |   | REFILLS                                  |
| Is this a first dose?   | Yes   | No If No, when was last dose given? _____   | When is patient due for next dose? _____ |
| ILUMYA  | 100mg SC injection at 0 and 4 weeks then every 12 weeks   |   | _____                                    |
| INFLIXIMAB<br>Avsola<br>Inlectra<br>Remicade<br>Renflexis   | <b>Induction:</b> _____ mg/kg or _____ mg IV infusion via gravity ---OR--- pump over at least 2 hours at weeks 0, 2, and 6  |   | NONE                                     |
|   | <b>Maintenance:</b> _____ mg/kg _____ mg IV infusion via gravity ---OR--- pump over at least 2 hours every _____ weeks<br>(Note: Round to nearest 100mg for Medicaid patients)  |   | _____                                    |
|   | If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.  |   | _____                                    |
| SIMPONI ARIA  | 2 mg/kg IV infusion via gravity ---OR--- pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter  |   | _____                                    |
| SPEVIGO   | 900 mg IV infusion over 90 minutes Additional 900 mg IV infusion over 90 minutes one week after initial dose if flare symptoms persist  |   | _____                                    |
| STELARA   | <b>Psoriasis Adult Subcutaneous</b><br>For patients <= 100 kg, 45 mg SC injection initially and 4 weeks later, followed by 45 mg every 12 weeks<br>For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks  |   | _____                                    |
|   | <b>Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose)</b><br>For patients <= 60 kg, 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks<br>For patients 60 kg – 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks<br>For patients >100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks |   | _____                                    |
|   | <b>Psoriatic Arthritis Adult</b><br>45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks<br>For patients with co-existent moderate-to-severe plaque psoriasis weighing > 100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks   |   | _____                                    |
| XOLAIR  | 150 or 300 mg SC injection once every 4 weeks   |   | _____                                    |
| IG  | <b>For Immunoglobulin therapy please refer to Immunoglobulin Form</b>   |   | _____                                    |
| OTHER   | _____   |   | _____                                    |
| <i>By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.</i> |   |   |  |

Prescriber's Signature \_\_\_\_\_  
Dispense as Written

Print Name

Date

ameritaiv.com

Prescriber's Signature \_\_\_\_\_  
Substitution Permitted

Print Name

Date

mosaiciv.com



ACHC ACCREDITED

