## Dermatology Order Form



Fax completed form to:

PATIENT INFORMATION						
Patient Name:	nt Name:		Date of Birth:		Referral Date:	
Address:		City/State/Zi		City/State/Zip:	<u>΄</u> ίρ:	
Home Phone:		Cell Phone:		Work Ph	none:	
Secondary Contact: He		Height:	Weight:	Male	e Female	
Patient Diagnosis & ICD-10:						
Allergies:						
PROVIDER INFORMATION						
Physician Name:		Lic.#:		DEA #:		
Practice Name:			NPI#:			
Address:			City/State/Zip:			
Office Contact: Phone:			Fax:			
Supervisory Physician (if applicable):						
PLEASE ATTACH						
Patient demographi	Patient demographics & front/back copy of all insurance cards (prescription & medical)  TB lab results within last 12 months (Stelara, Simponi Aria, Ilumya & Infliximabs only,					
Recent office visit notes, history & physical, lab & pertinent procedure results  HBV lab results within last 12 months (Infliximabs & Simponi Aria only)						
Current medication list & list of prior medications tried and failed (with dates)  Letter of medical necessity if drug dosing or indication is outside of FDA guidelines						
NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.						
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 10units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line						
Lab Orders: Lab Date & Frequency:						
PRESCRIPTION ORDERS						
Anaphylaxis Kit: Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed						
(Check all that apply) Diphenhydraminemq IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other						
Pre-Medications: Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IVminutes prior to infusion						
(Check all that apply) Diphenhydraminemg POOR IV infusionminutes prior to infusion Other						
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary						
PRODUCT			ION INFORMATI			REFILLS
Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose?						
ILUMYA	100mg SC injection at 0 and 4 weeks then eve	ry 12 weeks	-			
INFLIXIMAB	Induction:mg/kg or	mg IV infusion via	gravityOR pump or	ver at least 2 hours at w	reeks 0, 2, and 6	NONE
Avsola	Maintenance:mg/kg	mg IV infusion via	gravity OR pump o	ver at least 2 hours ever	ry weeks	
Inflectra	lectra (Note: Round to nearest 100mg for Medicaid patients)					
Remicade	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.					
Renflexis	<u> </u>			0 1 11 6		
SIMPONI ARIA	2 mg/kg IV infusion via gravity OR pump over 30 minutes at weeks 0 and 4, and every 8 weeks thereafter  900 mg IV infusion over 90 minutes Additional 900 mg IV infusion over 90 minutes one week after initial dose if flare symptoms persist					
SPEVIGO	-	Additional 900 mg IV infus	ion over 90 minutes one week	after initial dose if flare	e symptoms persist	
	Psoriasis Adult Subcutaneous	on initially and 4alsalsalat	on followed by 45 mar avenu 1	2aala		
STELARA	For patients <= 100 kg, 45 mg SC injection initially and 4 weeks later, followed by 45 mg every 12 weeks For patients > 100 kg, 90 mg SC injection initially and 4 weeks later, followed by 90 mg every 12 weeks					
	Psoriasis Pediatric Patients 6 to 17 (based on weight at time of initial dose)					
	For patients <= 60 kg, 0.75 mg/kg SC injection initially and 4 weeks later, then every 12 weeks					
	For patients 60 kg — 100kg, 45 mg SC injection initially and 4 weeks later, then every 12 weeks					
	For patients > 100kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks					
	Psoriatic Arthritis Adult					
	45 mg SC injection initially and 4 weeks later, followed by 45 mg SC injection every 12 weeks For patients with co-existent moderate-to-severe plaque psoriasis weighing >100 kg, 90 mg SC injection initially and 4 weeks later, then every 12 weeks					
XOLAIR	150 or 300 mg SC injection once every 4 weeks					
IG	For Immunoglobulin therapy please refer		m			
OTHER	10. minianogiocami diciupy picuse leiei					
By signing this form and utilizing our services, you are authorizing Eventus to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.						
, , , , , , , , , , , , , , , , , , ,						

Prescriber's Signature
Dispense as Written

**Print Name** 

Date

Prescriber's Signature Substitution Permitted **Print Name** 

Date





