Gastroenterology Order Form





Fax completed form to:

PATIENT INFORMATION						
Patient Name:		Date of Birth:		Refe	Referral Date:	
Address:				City/State/Zip:	ip:	
Home Phone:		Cell Phone:			k Phone:	
		Height:	Weight:	N	Male Female	
Patient Diagnosis & ICD-10:						
Allergies:						
PROVIDER INFORMATION						
,		Lic.#:	DEA #:			
Practice Name:			NPI#:			
Address:		DI.	City/State/Zip:			
		Phone: Fax:				
Supervisory Physician (if applicable): DLEASE ATTACH						
PLEASE ATTACH						
Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable Vaccine status (any vaccination) and documentation of any recent vaccinations TB lab results within last 12 months HBV lab results within last 12 months (Infliximabs only) Letter of medical necessity if drug dosing or indication is outside of FDA guidelines						
NURSING & LAB ORDERS						
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.						
Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line						
Lab Orders:						
Lab Date & Frequency:						
PRESCRIPTION ORDERS						
Anaphylaxis Kit:	Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed					
(Check all that apply)						
Pre-Medications:	Acetaminophenmg PO	minutes prior to i	nfusion Solu-Medi	rolmg IV	minutes prior to infusion	
(Check all that apply)	Diphenhydramine mg	PO OR IV	minutes prior to infusio	n Ot	ther	
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary						
PRODUCT		PRESCRIPTION	ON INFORMATION	ON		REFILLS
Is this a first dose?	es No If No, when was last dose given?	·	When is patient due for next d	lose?		
ENTYVIO	Induction : 300mg IV infusion via gr	ravityOR pump	over 30 minutes at week 0, 2,	and 6		NONE
	Maintenance: 300mg IV infusion via gravityOR pump over 30 minutes every weeks					
INFLIXIMAB Avsola	Induction:mg/kg or	mg IV infusion via	gravity 0R pump o	over at least 2 hours a	at weeks 0. 2. and 6	NONE
	Maintenance:mg/kg	ngtV infusion via				NONE
Inflectra	Maintenance:mg/kgmg/V infusion via gravity OR pump over at least 2 hours everyweeks(Note: Round to nearest 100mg for Medicaid patients)					
Remicade Renflexis If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.						
	Induction (Adult Dosing -Based on body weight of patient at time of dosing):					
STELARA	For patients 55kg or less administer 260mg IV infusion via gravityOR pump over at least 1 hour x 1 dose					
	For patients more than 55kg to 85kg administer 390mg IV infusion via gravity OR pump over at least 1 hour x 1 dose					NONE
	For patients more than 85kg administer 520mg IV infusion via gravityOR pump over at least 1 hour x 1 dose					NONL
	Maintenance: 90mg SubQ injection	weeks after ind	uction and every w	veeks thereafter		
OTHER						NONE
By signing this for	m and utilizing our services, you are authorizi	ng OptionOne to serve as y	our prior authorization design	nated agent in deali	ng with medical and prescription in:	surance companies.

<u>Dispense as Written</u>

Prescriber's Signature

Print Name

Date

Prescriber's Signature Substitution Permitted **Print Name**

Date





