Gastroenterology Order Form





Fax completed form to:

PATIENT INFORMATION					
Patient Name:		Date of Birth:		Referral Date:	
Address:				City/State/Zip:	
Home Phone:		Cell Phone:		Work Phone:	
Secondary Contact:		Height:	Weight:	Male Female	
Patient Diagnosis & ICD-	-10:				
Allergies:					
PROVIDER INFORMATION					
Physician Name:		Lic.#:		DEA #:	
Practice Name:			NPI#:		
Address:		City/State/Zip:			
		Phone: Fax:		Fax:	
Supervisory Physician (if applicable):					
PLEASE ATTACH					
Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicableVaccine status (any vaccination) and documentation of any recent vaccinatio TB lab results within last 12 months HBV lab results within last 12 months (Infliximabs only) Letter of medical necessity if drug dosing or indication is outside of FDA guide					
NURSING & LAB ORDERS					
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mLOR 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders:					
Lab Orders: Lab Date & Frequency:					
PRESCRIPTION ORDERS					
Anaphylaxis Kit:	Epinephrine 0.3mg IM as needed Solu-cortef 250mg-500mg IV as needed Solu-Medrol 60mg - 125mg IV as needed				
(Check all that apply)	Diphenhydramine mg IV as needed NS Hydration 500 ml IV over 30 minutes as needed Other				
Pre-Medications:	Acetaminophenmg POminutes prior to infusion Solu-Medrolmg IVminutes prior to infusion				
(Check all that apply) Diphenhydramine mg PO OR IV minutes prior to infusion Other					
Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary					
PRODUCT		PRESCRIPTI	ION INFORMATI	ON	REFILLS
Is this a first dose? Yes No If No, when was last dose given? When is patient due for next dose?					
ENTYVIO	Induction: 300mg IV infusion via	jravity 0R pump	o over 30 minutes at week 0, 2,	and 6	NONE
	Maintenance: 300mg IV infusion via gravity OR pump over 30 minutes every weeks				
INFLIXIMAB	Induction:mg/kg or	mg IV infusion via	gravity OR pump of	over at least 2 hours at weeks 0, 2, and 6	NONE
Avsola Inflectra	Maintenance:mg/kg	mgIV infusion via		ver at least 2 hours every weeks	
Remicade	(Note: Round to nearest 100mg for Medicaid patients)				
Renflexis	If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert.				
	Induction (Adult Dosing -Based on body weight of patient at time of dosing):				
STELARA	For patients 55kg or less administer 260mg IV infusion via gravityOR pump over at least 1 hour x 1 dose				
	For patients more than 55kg to 85kg administer 390mg IV infusion via gravityOR pump over at least 1 hour x 1 dose				NONE
	For patients more than 85kg administer 520mg IV infusion via gravityOR pump over at least 1 hour x 1 dose				
	Maintenance: 90mg SubQ injection weeks after induction and every weeks thereafter				
OTHER					NONE
By signing this form and utilizing our services, you are authorizing Amerita to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.					

Prescriber's Signature <u>Dispense as Written</u> Print Name

Date

Prescriber's Signature Substitution Permitted

Print Name

Date

©2022 Amerita Inc. All rights reserved. AME MOS GI ORDER FORM 06-09-22

ameritaiv.com

mosaiciv.com

