

# Multiple Sclerosis Order Form



Fax completed form to: \_\_\_\_\_

PATIENT INFORMATION			
Patient Name:	Date of Birth:	Referral Date:	
Address:		City/State/Zip:	
Home Phone:	Cell Phone:	Work Phone:	
Secondary Contact:	Height:	Weight:	Male Female
Patient Diagnosis & ICD-10:			
Allergies:			
PROVIDER INFORMATION			
Physician Name:	Lic.#:	DEA #:	
Practice Name:		NPI#:	
Address:		City/State/Zip:	
Office Contact:	Phone:	Fax:	
Supervisory Physician (if applicable):			
MS CLINICAL DETAILS			
<b>Type of MS:</b> Primary progressive multiple sclerosis (PPMS) ---OR--- Relapsing multiple sclerosis (RMS)			
<b>Ambulation status:</b> Able to ambulate more than 5 meters    Able to ambulate without aid or rest for at least 100 meters			
<b>Relapse details:</b> Two or more relapses within the previous two years    One relapse within the previous year			
PLEASE ATTACH			
Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable		Quantitative serum Immunoglobulin lab results ( <i>Ocrevus and Briumvi only</i> ) Vaccine status (any vaccination) and documentation of any recent vaccinations HBV lab results within last 12 months ( <i>Ocrevus and Briumvi only</i> ) Letter of medical necessity if drug dosing or indication is outside of FDA guideline	
NURSING & LAB ORDERS			
<b>Nurse Orders:</b> Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.			
<b>Flush Orders:</b> NaCl 0.9% - 5-10mL flush pre and post infusion and as needed    Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line			
<b>Lab Orders:</b> <span style="float: right;"><b>Lab Date &amp; Frequency:</b></span>			
PRESCRIPTION ORDERS			
<b>Anaphylaxis Kit:</b> (Check all that apply)	Epinephrine 0.3mg IM as needed Diphenhydramine _____ mg IV infusion as needed	Solu-cortef 250mg-500mg IV infusion as needed NS Hydration 500 ml IV infusion over 30 minutes as needed	Solu-Medrol 60mg - 125mg IV infusion as needed Other _____
<b>Pre-Medications:</b> (Check all that apply)	Acetaminophen _____ mg PO _____ minutes prior to infusion Diphenhydramine _____ mg PO ---OR--- IV infusion _____ minutes prior to infusion	Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion	Other _____
<b>Supply Orders:</b> All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary			
PRODUCT	PRESCRIPTION INFORMATION		REFILLS
Is this a first dose?	Yes	No	If No, when was last dose given? _____ When is patient due for next dose? _____
BRIUMVI	<b>Induction:</b> 150mg IV infusion via gravity ---OR--- pump over at least 4 hours followed 2 weeks later by 450mg IV infusion over at least 1 hour <b>Maintenance:</b> 450mg IV infusion via gravity ---OR--- pump over 1 hour 24 weeks after the first infusion and every 24 weeks thereafter <b>Post Infusion:</b> Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion (Per PI, Corticosteroid and antihistamine required for pre-medication, refer to section above)		NONE _____
OCREVUS	<b>Induction:</b> 300mg IV infusion via gravity ---OR--- pump over at least 2.5 hours followed 2 weeks later by 300mg IV infusion over at least 2.5 hours <b>Maintenance:</b> 600mg IV infusion via gravity ---OR--- pump over 3.5 hours every 6 months (if no prior serious infusion reactions, may administer over at least 2 hours) <b>Post Infusion:</b> Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion (Per PI, Corticosteroid and antihistamine required for pre-medication, refer to section above)		NONE _____
TYSABRI	300mg IV infusion via gravity ---OR--- pump over one hour every 4 weeks <b>Post Infusion:</b> Sodium Chloride 0.9% 100ml administer IV to keep line open (KVO) for one hour following infusion		NONE _____
IG	<b>For Immunoglobulin therapy please refer to Immunoglobulin Form</b>		
LEMTRADA	<b>For Lemtrada therapy please refer to Lemtrada Form</b>		
OTHER			_____
<b>By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.</b>			

Prescriber's Signature \_\_\_\_\_  
 Dispense as Written  
 Print Name \_\_\_\_\_ Date \_\_\_\_\_

Prescriber's Signature \_\_\_\_\_  
 Substitution Permitted  
 Print Name \_\_\_\_\_ Date \_\_\_\_\_



ACHC ACCREDITED

