Gastroenterology Referral Form





Fax Completed Form To: 877-418-4495 Phone: 877-418-4114

| | PATIENT INFORMATION | |
|---|--|-----------------------|
| Patient Name: | Date of Birth: Referral Date: | |
| Address: City/State/Zip: | | |
| Home Phone: | Cell Phone: Work Phone: | |
| Secondary Contact: | Height: Weight: □ Male □ Female | |
| Patient Diagnosis & ICD | -10: | |
| Allergies: | | |
| | PROVIDER INFORMATION | |
| Physician Name: | Lic.#: DEA#: | |
| Practice Name: NPI#: | | |
| Address: City/State/Zip: | | |
| Office Contact: | Phone: Fax: | |
| Supervisory Physician (it | fapplicable): | |
| | PLEASE ATTACH | |
| □ Patient demographics & front/back copy of all insurance cards (prescription & medical) □ TB lab results within last 12 months □ Recent office visit notes, history & physical, lab & pertinent procedure results □ HBV lab results within last 12 months (Infliximabs only) □ Current medication list & list of prior medications tried and failed (with dates) □ Liver enzymes lab results (Skyrizi only) □ Line access documentation/verification if applicable □ Bilirubin levels (Skyrizi only) □ Vaccine status (any vaccination) and documentation of any recent vaccinations □ Letter of medical necessity if drug dosing or indication is outside of FDA guidelines | | s |
| | NURSING & LAB ORDERS | |
| | o provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. % - 5-10mL flush pre and post infusion and as needed Heparin - 100units/mL | ated to maintain line |
| | PRESCRIPTION ORDERS | |
| Anaphylaxis Kit: (Check all that apply) | ☐ Epinephrine 0.3mg IM as needed ☐ Solu-cortef 250mg-500mg IV as needed ☐ Solu-Medrol 60mg - 12☐ Diphenhydramine mg IV as needed ☐ NS Hydration 500 ml IV over 30 minutes as needed ☐ Other | 25mg IV as needed |
| Pre-Medications: (Check all that apply) | ☐ Acetaminophenmg PO minutes prior to infusion ☐ Solu-Medrolmg IVminutes prior to infusion ☐ Diphenhydraminemg ☐ POOR ☐ IVminutes prior to infusion ☐ Other | |
| Supply Orders: All sup | plies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary | |
| PRODUCT | PRESCRIPTION INFORMATION | REFILLS |
| Is this a first dose? | Yes 🔲 No If No, when was last dose given?When is patient due for next dose? | |
| □ ENTYVIO | ☐ Induction: 300mg IV infusion over 30 minutes at week 0, 2, and 6 | NONE |
| | ☐ Maintenance: 300mg IV infusion over 30 minutes every weeks weeks weeks 6 | 2 pens, 13 refills |
| ☐ INFLIXIMAB ☐ Avsola ☐ Inflectra ☐ Remicade | □ Induction:mg/kg ormg IV infusion via □ gravityOR □ pump over at least 2 hours at weeks 0, 2, and 6 | NONE |
| | ☐ Maintenance :mg/kgmg/V infusion via ☐ gravity OR ☐ pump over at least 2 hours everyweeks (Note: Round to nearest 100mg for Medicaid patients) | |
| Renflexis | If Remicade infusion tolerated, adjust infusion time according to manufacturer package insert. | |
| □ ом∨он | ☐ Induction: 300mg IV infusion via ☐ gravityOR ☐ pump over 30 minutes at week 0, 4, and 8 | NONE |
| | Maintenance: 200mg SC injection (given as two consecutive injections of 100 mg each) at Week 12, and every 4 weeks thereafter | |
| □ SKYRIZI | ☐ Induction (Crohn's): 600mg IV infusion via ☐ gravityOR ☐ pump over one hour at week 0, 4, and 8 ☐ Maintenance: ☐ 180mg or ☐ 360mg SC injection at Week 12, and every 8 weeks thereafter | NONE |
| | □ Induction (UC): 1200mg IV infusion via □ gravityOR□ pump over two hours at week 0, 4, and 8 □ Maintenance: □ 180mg or □ 360mg SC injection at Week 12, and every 8 weeks thereafter | NONE |
| ☐ STELARA | Induction (Adult Dosing -Based on body weight of patient at time of dosing): □ For patients 55kg or less administer 260mg IV infusion via □ gravityOR□ pump over at least 1 hour x 1 dose □ For patients more than 55kg to 85kg administer 390mg IV infusion via □ gravityOR□ pump over at least 1 hour x 1 dose □ For patients more than 85kg administer 520mg IV infusion via □ gravityOR□ pump over at least 1 hour x 1 dose □ Hainton and Source Sour | NONE |
| | ☐ Maintenance: 90mg SubQ injectionweeks after induction and everyweeks thereafter | |
| | ☐ Induction: 200mg IV infusion on weeks 0, 4, and 8 | NONE |
| ☐ TREMFYA | | NONE |
| ☐ TREMFYA | ☐ Induction: 200mg IV infusion on weeks 0, 4, and 8 ☐ Maintenance: 100mg SubQ injection every 8 weeks beginning at week 16 | NONE |

Prescriber's Signature <u>Dispense as Written</u>

Print Name

Date

Prescriber's Signature Substitution Permitted

Print Name

Date



