

# Ultomiris Order Form



Fax completed form to: \_\_\_\_\_

## PATIENT INFORMATION

Patient Name:		Date of Birth:		Referral Date:	
Address:			City/State/Zip:		
Home Phone:		Cell Phone:		Work Phone:	
Secondary Contact:		Height:                      Weight:		Male      Female	
Patient Diagnosis & ICD-10:					
Allergies:					

## PROVIDER INFORMATION

Physician Name:		Lic.#:		DEA #:	
Practice Name:			NPI#:		
Address:					
Office Contact:		Phone:		Fax:	
Supervisory Physician (if applicable):					

## PLEASE ATTACH

Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable	Vaccine status (any vaccination) and documentation of any recent vaccinations Clinical documentation on any recent meningococcal infections Documentation of a meningococcal vaccination Letter of medical necessity if drug dosing or indication is outside of FDA guidelines
---	---

## NURSING & LAB ORDERS

**Nurse Orders:** Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders.

**Flush Orders:** NaCl 0.9% - 5-10mL flush pre and post infusion and as needed    Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line

**Lab Orders:** **Lab Date & Frequency:**

## PRESCRIPTION ORDERS

<b>Anaphylaxis Kit:</b>	Epinephrine 0.3mg IM as needed	Solu-cortef 250mg-500mg IV infusion as needed	Solu-Medrol 60mg - 125mg IV infusion as needed
(Check all that apply)	Diphenhydramine _____ mg IV infusion as needed	NS Hydration 500 ml IV infusion over 30 minutes as needed	Other
<b>Pre-Medications:</b>	Acetaminophen _____ mg PO _____ minutes prior to infusion	Solu-Medrol _____ mg IV infusion _____ minutes prior to infusion	
(Check all that apply)	Diphenhydramine _____ mg PO ---OR--- IV infusion _____ minutes prior to infusion	Other	

**Supply Orders:** All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
---------	--------------------------	---------

Is this a first dose?    Yes    No    If No, when was last dose given? \_\_\_\_\_    When is patient due for next dose? \_\_\_\_\_

Is the prescriber enrolled in the Ultomiris REMS program?    Yes    No

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Ultomiris  PNH and aHUS	<b>Loading Dose</b> For patients 5-10kg administer 600mg IV infusion via gravity ---OR--- pump over at least 1.4 hours For patients 10-20kg administer 600mg IV infusion via gravity ---OR--- pump over at least 0.8 hours For patients 20-30kg administer 900mg IV infusion via gravity ---OR--- pump over at least 0.6 hours For patients 30-40kg administer 1,200mg IV infusion via gravity ---OR--- pump over at least 0.5 hours	NONE
	For patients 40-60kg administer 2,400mg IV infusion via gravity ---OR--- pump over at least 0.8 hours For patients 60-100kg administer 2,700mg IV infusion via gravity ---OR--- pump over at least 0.6 hours For patients >100kg administer 3,000mg IV infusion via gravity ---OR--- pump over at least 0.4 hours	
PNH, aHUS and gMG	<b>Maintenance Dose</b> For patients 5-10kg administer 300mg IV infusion via gravity ---OR--- pump over at least 0.8 hours every 4 weeks For patients 10-20kg administer 600mg IV infusion via gravity ---OR--- pump over at least 0.8 hours every 4 weeks For patients 20-30kg administer 2,100 IV infusion via gravity ---OR--- pump over at least 1.3 hours every 8 weeks For patients 30-40kg administer 2,700mg IV infusion via gravity ---OR--- pump over at least 1.1 hours every 8 weeks	_____
	For patients 40-60kg administer 3,000mg IV infusion via gravity ---OR--- pump over at least 0.9 hours every 8 weeks For patients 60-100kg administer 3,300mg IV infusion via gravity ---OR--- pump over at least 0.7 hours every 8 weeks For patients >100kg administer 3,600mg IV infusion via gravity ---OR--- pump over at least 0.5 hours every 8 weeks	
OTHER		NONE

*By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.*

Prescriber's Signature \_\_\_\_\_    Print Name \_\_\_\_\_    Date \_\_\_\_\_  
**Dispense as Written**

Prescriber's Signature \_\_\_\_\_    Print Name \_\_\_\_\_    Date \_\_\_\_\_  
**Substitution Permitted**

